

SHIN'ICHI NOJIRI

Department of Mathematics and Physics

National Defense Academy

Hashirimizu, Yokosuka 239, JAPAN

ICHIRO ODA

NDA-FP-5/92, OCHA-PP-27
July 1992

Semi-classical Approach to Charged
Dilatonic Black Hole in Two Dimensions

SHIN'ICHI NOJIRI

Department of Mathematics and Physics

National Defense Academy

Hashirimizu, Yokosuka 239, JAPAN

ICHIRO ODA

Faculty of Science, Department of Physics

Ochanomizu University

1-1 Otsuka 2, Bunkyo-ku, Tokyo 112, JAPAN

ABSTRACT

We consider exactly solvable semi-classical theory of two dimensional dilatonic gravity with electromagnetic interactions. As was done in the paper by Russo, Susskind and Thorlacius, the term which changes the kinetic term is added to the action. The theory contains massless fermions as matter fields and there appear the quantum corrections including chiral anomaly. The screening effect due to the chiral anomaly has a tendency to cloak the singularity. In a region of the parameter space, the essential behavior of the theory is similar to that of Callan, Giddings, Harvey and Strominger's dilatonic black hole theory modified in the paper by Russo, Susskind and Thorlacius and the singularity formed by the collapsing matter emerges naked. We find, however, another region of the parameter space where the singularity disappears in a finite proper time. Furthermore, in the region of the parameter space, there appears a discontinuity in the metric on the trajectory of the collapsing matter, which would be a signal of topology change.

1. Introduction

One of most important theme in elementary particle physics is to construct the quantum theory of the gravity. Black hole evaporation, however, provides a serious problem. A quantum mechanically pure state which describes gravitationally collapsing matter to form a black hole, evolves into a mixed quantum state which describes Hawking radiation.¹ This implies that there does not exist a well-defined S-matrix theory. There are several scenarios to solve this problem of the loss of quantum informations. Hawking has given a most radical proposal that the black hole completely evaporates and the quantum coherence is lost in the gravitational collapse. On the other hand, 't Hooft has proposed that Hawking radiation carries off informations about the quantum states of the black holes.² Another conservative proposal is that the process of the collapse and the radiation leaves a stable remnant which carries the information of the initial configuration of the system.³ However, if the remnant has the mass of order the Planck scale, gravitational effects would produce the light remnants in pair and the lifetime of stars would be shorter than observed. Therefore the mass of the remnants should be macroscopic. A candidate of such remnants is the extremal black holes of Reissner-Nordström solutions.⁴ The extremal black holes have vanishing temperatures and the space-time metric is not singular everywhere.

Recently, Callan, Giddings, Harvey and Strominger (CGHS) have investigated an interesting toy model of two dimensional gravity.⁵ In this model, the gravity couples with a dilaton and conformal matter fields. They found classical exact solutions, including the solutions describing the formation of a black hole by collapsing matter. Furthermore Hawking radiation and the back reaction of the metric can be described by adding correction terms to the classical equations of motion. The correction terms derive from the conformal anomaly of the matter fields. It was also argued that the quantum correction would remove the usual black hole singularity. Several groups,⁶⁻⁹ however, pointed that gravitational collapse always leads to a curvature singularity. Susskind and Thorlacius have found massive static

solutions with zero temperature but they have shown that a decaying black hole cannot evolve into one of them.⁷

Although the original CGHS semi-classical equations of motion have not been solved explicitly, Bilal and Callan¹⁰ and de Alwis¹¹ have found that the original model can be modified to allow obtaining explicit exact solutions of quantum black holes.^{*} After that, Russo, Susskind and Thorlacius have studied a variation.¹³ They have changed a kinetic term by adding the following term to the action,

$$\delta S = -\frac{\kappa}{4\pi} \int d^2x \sqrt{-g} \phi R . \quad (1.1)$$

Here ϕ is a dilaton field and R is a scalar curvature. Then there appears a conserved current J^μ , which has the following form,

$$J^\mu = \partial^\mu(\phi - \rho) , \quad (1.2)$$

in the conformal gauge,

$$g_{\mp\pm} = -\frac{1}{2}e^{2\rho} , \quad g_{\pm\pm} = 0 . \quad (1.3)$$

The current has already appeared in CGHS's classical theory. The conservation law $\partial_\mu J^\mu = \partial_\mu \partial^\mu(\phi - \rho) = 0$ allows to fix the residual gauge symmetry of reparametrization by choosing the condition $\phi - \rho = 0$, which is given by a sum of holomorphic function $w^+(x^+)$ and anti-holomorphic function $w^-(x^-)$ in general : $\phi(x^+, x^-) - \rho(x^+, x^-) = w^+(x^+) + w^-(x^-)$. The gauge choice is very useful when we construct explicit solutions of classical and semi-classical equations of motion.

In the previous paper,¹⁴ the authors have coupled electromagnetic fields to CGHS dilatonic gravity. The coupling has been chosen so that the current (1.2) is conserved. This model can be also solved classically. Among the solutions describing static black holes, there exist extremal solutions which have zero temperatures.

^{*} Similar but different quantum model was proposed by Hamada.¹²

In the extremal solutions, the space-time metric is not singular and these solutions should be analogues of four dimensional extremal Reissner-Nordström black holes. We also obtain the solutions describing charged matters (massless fermions) collapsing into black holes. Through the collapse, not only future horizon but past horizon is also shifted. The quantum corrections have been also discussed by including chiral anomaly. When the matter collapses into a black hole, except the case that the resulting black hole is extremal, the curvature singularity also appeared in the quantum theory in a way similar to CGHS model.

In this paper, we consider exactly solvable semi-classical theory of two dimensional dilatonic gravity with electromagnetic interactions. The term corresponding to (1.1) is added to the action. The screening effect due to the chiral anomaly has a tendency to cloak the singularity. In a region of the parameter space, the behavior of the theory is similar to that of modified CGHS's dilatonic black hole theory which has been analyzed by Russo, Susskind and Thorlacius¹³ and the singularity formed by the collapsing matter emerges naked. In another region of the parameter space, however, the singularity disappears in a finite proper time. Furthermore, in the region of the parameter space, there appears a discontinuity in the metric on the trajectory of the collapsing matter, which would be a signal of topology change.

In the next section, we review the classical theory of charged dilatonic black holes based on the previous paper. In section 3, we consider the semi-classical theory by adding the correction terms to the classical action. The correction terms contain the terms which come from the conformal and chiral anomalies and the term corresponding to (1.1). The last section is devoted to summary and discussion.

2. Classical Theory of Charged Dilatonic Black Holes

We start with the following action,

$$S_c = \frac{1}{2\pi} \int d^2x \sqrt{-g} \{ e^{-2\phi} (R + 4(\nabla\phi)^2 + 4\lambda^2) - \frac{e^{a\phi}}{g_A^2} F^2 - \sum_{j=1}^N i \bar{\Psi}_j \gamma^\mu (D_\mu - iA_\mu) \Psi_j \} . \quad (2.1)$$

Here $\Psi_j = \begin{pmatrix} \psi_j^L \\ \psi_j^R \end{pmatrix}$'s are N massless complex fermions. g_A is a $U(1)$ electromagnetic gauge coupling constant and D_μ is a covariant derivative. If we fix $a = -2$, this action, except the fermion part, describes an effective field theory derived from string theory.^{15,16} In this paper, we choose $a = 2$, which gives a conserved current (1.2) and we can obtain the classical exact solutions of this model. When we fix the gauge freedom of reparametrization invariance by the conformal gauge (1.3) and that of $U(1)$ gauge symmetry by the following light-cone gauge condition

$$A_- = 0 , \quad (2.2)$$

the action (2.1) reduces into

$$S_c = \frac{1}{2\pi} \int d^2x \{ e^{-2\phi} (4\partial_+ \partial_- \rho - 8\partial_+ \phi \partial_- \phi + 2\lambda^2 e^{2\rho}) + \frac{4}{g_A^2} e^{2\phi-2\rho} F_{+-}^2 + \frac{i}{2} \sum_{j=1}^N (\psi_j^{L*} \partial_- \psi_j^L + \psi_j^{R*} (\partial_+ - iA_+) \psi_j^R) \} . \quad (2.3)$$

Here $F_{+-} = -\partial_- A_+$. The metric equations are given by

$$0 = T_{++} = e^{-2\phi} (4\partial_+ \rho \partial_+ \phi - 2\partial_+^2 \phi) + \frac{i}{4} \sum_{j=1}^N (\psi_j^{L*} \partial_+ \psi_j^L - \partial_+ \psi_j^{L*} \psi_j^L) + \frac{1}{2} A_+ \sum_{j=1}^N \psi_j^{L*} \psi_j^L , \quad (2.4)$$

$$0 = T_{--} = e^{-2\phi} (4\partial_- \rho \partial_- \phi - 2\partial_-^2 \phi) + \frac{i}{4} \sum_{j=1}^N (\psi_j^{R*} \partial_- \psi_j^R - \partial_- \psi_j^{R*} \psi_j^R) ,$$

$$0 = T_{+-} = e^{-2\phi}(2\partial_+\partial_-\phi - 4\partial_+\phi\partial_-\phi - \lambda^2 e^{2\rho}) + \frac{2}{g_A^2}e^{2\phi-2\rho}F_{+-}^2 . \quad (2.5)$$

The Maxwell equations are

$$\begin{aligned} 0 &= -\frac{8}{g_A^2}\partial_+(e^{2(\phi-\rho)}F_{+-}) + \frac{1}{2}\sum_{j=1}^N\psi_j^{L*}\psi_j^L , \\ 0 &= \frac{8}{g_A^2}\partial_-(e^{2(\phi-\rho)}F_{+-}) + \frac{1}{2}\sum_{j=1}^N\psi_j^{R*}\psi_j^R . \end{aligned} \quad (2.6)$$

We also obtain dilaton and matter equations of motion,

$$0 = -4\partial_+\partial_-\phi + 4\partial_+\phi\partial_-\phi + 2\partial_+\partial_-\rho + \lambda^2 e^{2\rho} - \frac{2}{g_A^2}e^{4\phi-2\rho}F_{+-}^2 , \quad (2.7)$$

$$0 = \partial_-\psi_j^L , \quad 0 = (\partial_+ - iA_+)\psi_j^R . \quad (2.8)$$

By using Equations (2.5) and (2.7), we find that the current J^μ defined in Equation (1.2) is conserved. Therefore we can fix the residual gauge symmetry of reparametrization by choosing the condition, $\phi - \rho = 0$. In the following, we only consider the solutions with $\psi_j^R = 0$. Then the general solutions of Equations (2.4) - (2.8) are given by,

$$\begin{aligned} \psi_j^L &= \psi_j^L(x^+) , \\ A_+ &= -\frac{g_A^2}{4}x^- \int^{x^+} dy^+ \sum_{j=1}^N \psi_j^{L*}(y^+)\psi_j^L(y^+) , \\ e^{-2\phi} = e^{-2\rho} &= \frac{M}{\lambda} - \lambda^2 x^+ x^- + \frac{g_A^2}{8}x^- \int^{x^+} dy^+ \left(\int^{y^+} dz^+ \sum_{j=1}^N \psi_j^{L*}(z^+)\psi_j^L(z^+) \right)^2 \\ &\quad - \frac{i}{4} \int^{x^+} dy^+ \int^{y^+} dz^+ \sum_{j=1}^N (\psi_j^{L*}(z^+)\partial_+\psi_j^L(z^+) - \partial_+\psi_j^{L*}(z^+)\psi_j^L(z^+)) . \end{aligned} \quad (2.9)$$

Now we consider solutions with $\psi_j^L = 0$. A special solution describing a dilaton

vacuum is given by,

$$A_+ = 0, \quad e^{-2\phi} = e^{-2\rho} = -\lambda^2 x^+ x^- . \quad (2.10)$$

We have more general solutions,

$$\begin{aligned} F_{+-} &= C \quad (\text{constant}) . \\ e^{-2\phi} &= e^{-2\rho} = \frac{M}{\lambda} - \tilde{\lambda}^2 x^+ x^- . \end{aligned} \quad (2.11)$$

Here $\frac{M}{\lambda}$ is an integration constant and $\tilde{\lambda}$ is defined by

$$\tilde{\lambda}^2 \equiv \lambda^2 - \frac{2C^2}{g_A^2} . \quad (2.12)$$

The solutions (2.11) tell that the metric has a singularity when $\frac{M}{\lambda} - \tilde{\lambda}^2 x^+ x^- = 0$ and there appear horizons $x^+ x^- = 0$. Therefore, if $C \neq 0$, the solutions (2.11) describe charged black holes. The structure of space-time is similar to that of the Schwarzschild black holes and simple in contrast to the Reissner-Nordström black hole solutions in four dimensions. Note that the singularity vanishes when $\tilde{\lambda}^2 = 0$ when we fix M to be finite, although the solution corresponds to a massive[★] and charged object. This solution could be a natural analogue of extremal Reissner-Nordström black hole solution. In fact this solution has a vanishing temperature. The temperature can be found by changing the coordinates

$$\begin{aligned} x^+ &= \frac{1}{\tilde{\lambda}} \sqrt{\frac{M}{\lambda}} e^{\tilde{\lambda} t} \sinh \tilde{\lambda} r \\ x^- &= -\frac{1}{\tilde{\lambda}} \sqrt{\frac{M}{\lambda}} e^{-\tilde{\lambda} t} \sinh \tilde{\lambda} r \end{aligned} \quad (2.13)$$

and Wick rotating $t \longrightarrow i\tau$. The resulting metric is given by

$$ds^2 = dr^2 + \tanh^2 \tilde{\lambda} r \, d\tau^2 . \quad (2.14)$$

The geometry approaches to Euclidean flat space-time when r goes to ∞ , $ds^2 \sim$

★ Here “massive” means that $\frac{M}{\lambda}$ does not vanish.

$dr^2 + d\tau^2$. On the other hand, the metric has a following form when $r \rightarrow 0$,

$$ds^2 \sim dr^2 + \tilde{\lambda}^2 r^2 d\tau^2 . \quad (2.15)$$

Since the metric is not singular if and only if τ has a period of $\frac{2\pi}{\tilde{\lambda}}$, the temperature T is given by,

$$T = \frac{\tilde{\lambda}}{2\pi} . \quad (2.16)$$

This equation (2.16) tells that the extremal solutions, which correspond to $\tilde{\lambda}^2 = 0$, have a vanishing temperature. This result should compare with CGHS black holes, which have a common non-vanishing temperature $\frac{\lambda}{2\pi}$. When $\tilde{\lambda}^2 < 0$, the temperature is imaginary and the naked singularity appears.

If we fix $\tilde{M} = \tilde{\lambda}^2 M$ to be finite and redefine the coordinates $x^\pm$ by,

$$x^+ \rightarrow x^+ + \frac{\alpha \tilde{M}}{\lambda \tilde{\lambda}^2} , \quad x^- \rightarrow x^- + \beta + \frac{1}{\alpha \tilde{\lambda}^2} , \quad (2.17)$$

we obtain the following metric, in the limit of $\tilde{\lambda}^2 \rightarrow 0$,

$$e^{-2\phi} = e^{-2\rho} = \frac{1}{\alpha} x^+ + \alpha \lambda \tilde{M} x^- + \beta . \quad (2.18)$$

The curvature is given by,

$$\begin{aligned} R &= 8e^{-2\rho} \partial_+ \partial_- \rho \\ &= \frac{4\lambda \tilde{M}}{\frac{1}{\alpha} x^+ + \alpha \lambda \tilde{M} x^- + \beta} , \end{aligned} \quad (2.19)$$

and we find that there appears a singularity when $\frac{1}{\alpha} x^+ + \alpha \lambda \tilde{M} x^- + \beta = 0$.

The solutions (2.9) can also describe charged matter (chiral fermions) collapsing into black holes. For example, we can consider a charged shock wave which is given

by,

$$\begin{aligned} \frac{i}{4} \sum_{j=1}^N (\psi_j^{L*}(x^+) \partial_+ \psi_j^L(x^+) - \partial_+ \psi_j^{L*}(x^+) \psi_j^L(x^+)) &= a \delta(x^+ - x_0^+) , \\ \frac{g_A^2}{4} \sum_{j=1}^N \psi_j^{L*}(x^+) \psi_j^L(x^+) &= b \delta(x^+ - x_0^+) . \end{aligned} \quad (2.20)$$

Then the solution is given by,

$$\begin{aligned} x^+ < x_0^+ , \quad F_{+-} &= C , \\ e^{-2\phi} &= e^{-2\rho} = \frac{M}{\lambda} - (\lambda^2 - \frac{2C^2}{g_A^2}) x^+ x^- , \\ x^+ > x_0^+ , \quad F_{+-} &= C + b , \\ e^{-2\phi} &= e^{-2\rho} = \frac{M}{\lambda} + a x_0^+ + \frac{2B a x_0^+}{g_A^2 \tilde{\lambda}'^2} \\ &\quad - \tilde{\lambda}'^2 (x^+ + \frac{2B x_0^+}{g_A^2 \tilde{\lambda}'^2}) (x^- + \frac{a}{\tilde{\lambda}'^2}) . \end{aligned} \quad (2.21)$$

Here $\tilde{\lambda}'$ and B are defined by,

$$\tilde{\lambda}'^2 \equiv \lambda^2 - \frac{2(C+b)^2}{g_A^2} , \quad B \equiv (2Cb + b^2) . \quad (2.22)$$

When $\tilde{\lambda}' = 0$, the metric (2.21) in case $x^+ > x_0^+$ corresponds to the metric in Equation (2.18). Note that there appears a singularity even in case $\tilde{\lambda}' = 0$.

Equation (2.21) tells that the event horizon when $x^+ > x_0^+$ is given by

$$(x^+ + \frac{2B x_0^+}{g_A^2 \tilde{\lambda}'^2}) (x^- + \frac{a}{\tilde{\lambda}'^2}) = 0 . \quad (2.23)$$

Note that not only future horizon corresponding to $x^- = -\frac{a}{\tilde{\lambda}'^2}$, but past horizon, where $x^+ = -\frac{2B x_0^+}{g_A^2 \tilde{\lambda}'^2}$, is shifted by the shock wave. In the original paper by 't Hooft,² the shift of only future horizon was discussed in four dimensional black holes.

The shift of the past horizon, which is observed in this paper, would suggest that charged particles collapsing into Reissner-Nordström black hole in four dimensions could shift the past horizon. The incoming matters cause the change of the charge distribution, which would shift the past horizon.

3. Semi-classical Analysis of Hawking Evaporation

In the original paper by CGHS, they have found that the Hawking radiation and the back reaction of the metric can be described by adding correction terms to the classical equation of motion. The correction terms come from the conformal anomaly. Now, since we have massless fermions, the terms which come from the chiral anomaly should be also added. Furthermore, by following to Russo, Susskind and Thorlacius,¹³ we also add a term S_R corresponding to δS in Equation (1.1), to preserve the conservation of the current J^μ in Equation (1.2). Then the total quantum action S_q is given by,

$$\begin{aligned}
S_q &= S_c + S_\rho + S_\chi + S_R , \\
S_\rho &= \frac{1}{2\pi} \int d^2x \sqrt{-g} \left\{ -\frac{1}{2}(\nabla Z)^2 + \sqrt{\frac{N}{48}} Z R \right\} , \\
S_\chi &= \frac{1}{2\pi} \int d^2x \left\{ \frac{1}{2} \sqrt{-g} (\nabla Y)^2 + \sqrt{\frac{N}{2}} Y \epsilon^{\mu\nu} F_{\mu\nu} \right\} , \\
S_R &= -\frac{N}{96\pi} \int d^2x \sqrt{-g} \phi R .
\end{aligned} \tag{3.1}$$

Here S_c is a classical action in Eq.(2.1). By adding new degrees of freedom Z and Y , we write the actions in local forms. We have neglected the one loop and higher order contributions from the reparametrization ghosts, Liouville mode and dilaton field by assuming that the number of fermions N is very large.

In the following, we only consider the solutions where fermions $\psi_j^{L,R}$ vanish. By choosing the gauge fixing conditions (1.3) and (2.2), the classical equations of

motion (2.4) - (2.6) are modified as follows,

$$\begin{aligned}
0 = T_{++} &= (e^{-2\phi} + \frac{N}{96})(4\partial_+\rho\partial_+\phi - 2\partial_+^2\phi) + \frac{1}{2}\partial_+Z\partial_+Z - \sqrt{\frac{N}{12}}\partial_+\rho\partial_+Z \\
&\quad + \frac{1}{2}\sqrt{\frac{N}{12}}\partial_+^2Z - \frac{1}{2}\partial_+Y\partial_+Y, \\
0 = T_{--} &= (e^{-2\phi} + \frac{N}{96})(4\partial_-\rho\partial_-\phi - 2\partial_-^2\phi) + \frac{1}{2}\partial_-Z\partial_-Z - \sqrt{\frac{N}{12}}\partial_-\rho\partial_-Z \\
&\quad + \frac{1}{2}\sqrt{\frac{N}{12}}\partial_-^2Z - \frac{1}{2}\partial_-Y\partial_-Y,
\end{aligned} \tag{3.2}$$

$$\begin{aligned}
0 = T_{+-} &= e^{-2\phi}(2\partial_+\partial_-\phi - 4\partial_+\phi\partial_-\phi - \lambda^2 e^{2\rho}) + \frac{2}{g_A^2}e^{2\phi-2\rho}F_{+-}^2 \\
&\quad - \sqrt{\frac{N}{48}}\partial_+\partial_-Z + \frac{N}{48}\partial_+\partial_-\phi,
\end{aligned} \tag{3.3}$$

$$\begin{aligned}
0 &= -\frac{8}{g_A^2}\partial_+(e^{2(\phi-\rho)}F_{+-}) - \sqrt{2N}\partial_+Y, \\
0 &= -\frac{8}{g_A^2}\partial_-(e^{2(\phi-\rho)}F_{+-}) - \sqrt{2N}\partial_-Y.
\end{aligned} \tag{3.4}$$

The dilaton equation of motion (2.7) is modified to be,

$$\begin{aligned}
0 &= -4\partial_+\partial_-\phi + 4\partial_+\phi\partial_-\phi + 2\partial_+\partial_-\rho + \lambda^2 e^{2\rho} \\
&\quad - \frac{2}{g_A^2}e^{4\phi-2\rho}F_{+-}^2 + e^{2\phi}\frac{N}{48}\partial_+\partial_-\rho,
\end{aligned} \tag{3.5}$$

We also have Z and Y equations,

$$0 = -2\partial_+\partial_-Z + \sqrt{\frac{N}{3}}\partial_+\partial_-\rho, \tag{3.6}$$

$$0 = 2\partial_+\partial_-Y + \sqrt{2N}F_{+-}. \tag{3.7}$$

By using Equations (3.3) and (3.5), we find $\partial_\mu\partial^\mu(\phi-\rho) = 0$ and the current J^μ defined in (1.2) is conserved. Therefore we can impose the residual gauge condition

$\rho = \phi$. Under this gauge condition, the modified Maxwell equation (3.4) can be solved with respect to Y as follows

$$\sqrt{2N}Y = -\frac{8}{g_A^2}F_{+-} + c . \quad (3.8)$$

Here c is a constant. By substituting this equation (3.8) into Equation (3.7), we obtain,

$$0 = \partial_+ \partial_- F_{+-} - \frac{g_A^2 N}{8} F_{+-} . \quad (3.9)$$

This equation tells that electromagnetic fields become massive due to the screening effect by fermion loops. If we consider a situation where an incoming shock wave (2.20) carries charge, the boundary are given by,

$$\begin{aligned} F_{+-} &= C \quad \text{when } x^+ = x_0^+ , \\ \partial_+ F_{+-} &= 0 \quad \text{when } x^+ = x_0^+ \text{ and } x^- = \tilde{x}_0 . \end{aligned} \quad (3.10)$$

The latter condition is imposed to fix the constant of integration.[★] The solution of Equation (3.9) with the boundary condition (3.10) is given by,

$$\begin{aligned} F_{+-} &= C J_0(p) \quad (x^- < \tilde{x}_0) \\ &= C I_0(q) \quad (x^- > \tilde{x}_0) . \end{aligned} \quad (3.11)$$

Here J_n is a Bessel function and I_n is a modified Bessel function:

$$\begin{aligned} J_n(z) &= \left(\frac{z}{2}\right)^n \sum_{m=0}^{\infty} \frac{(-1)^m (z/2)^{2m}}{m! \Gamma(m+n+1)} , \\ I_n(z) &= \left(\frac{z}{2}\right)^n \sum_{m=0}^{\infty} \frac{(z/2)^{2m}}{m! \Gamma(m+n+1)} , \end{aligned} \quad (3.12)$$

★ Since the velocity of massless matter is larger than that of electro-magnetic fields, which is massive, the trajectory of the matter makes a shock wave of electro-magnetic field like Cherenkov radiation. The parameter $\tilde{x}_0$ is related to the strength of the shock wave.

and p and q are defined by,

$$p \equiv \sqrt{\frac{g_A^2 N (\tilde{x}_0 - x^-)(x^+ - x_0^+)}{2}}, \quad q \equiv \sqrt{-\frac{g_A^2 N (\tilde{x}_0 - x^-)(x^+ - x_0^+)}{2}}. \quad (3.13)$$

This screening effect makes the field strength F_{+-} decrease when $x^- < \tilde{x}_0$ *i.e.*, in the asymptotic region and increase near when $x^- > \tilde{x}_0$. As we will see later, the singularity can disappear in a finite proper time due to this screening effect.

Equation (3.6) can be also solved with respect to Z as follows,

$$Z = \sqrt{\frac{N}{12}} \rho + r^+(x^+) + r^-(x^-). \quad (3.14)$$

By using this equation and the gauge condition $\rho = \phi$, Equation (3.2) can be written as follows,

$$\begin{aligned} 0 &= \partial_+^2 (e^{-2\phi} + \frac{N}{48} \phi) + t^+(x^+) - \frac{1}{4N} \left\{ \partial_+ \left(-\frac{8}{g_A^2} F_{+-} \right) \right\}^2 \\ 0 &= \partial_-^2 (e^{-2\phi} + \frac{N}{48} \phi) + t^-(x^-) - \frac{1}{4N} \left\{ \partial_- \left(-\frac{8}{g_A^2} F_{+-} \right) \right\}^2. \end{aligned} \quad (3.15)$$

Here $t^\pm(x^\pm)$ is defined by

$$t^\pm(x^\pm) \equiv \frac{1}{2} (\partial_\pm r^\pm)^2 + \frac{1}{4} \sqrt{\frac{N}{3}} \partial_\pm^2 r^\pm. \quad (3.16)$$

Since $r^\pm$ is a solution of the equation motion (3.6), $t^\pm(x^\pm)$ represents dynamical degrees of freedom. Furthermore we obtain the following equations by using Equations (3.3), (3.5), and (3.14) and the gauge condition $\phi - \rho = 0$,

$$0 = \partial_+ \partial_- (e^{-2\phi} + \frac{N}{48} \phi) + \lambda^2 - \frac{2}{g_A^2} F_{+-}^2. \quad (3.17)$$

The classical solution describing the dilaton vacuum (2.10) satisfies Equations (3.9) and (3.17). If we require that the solution also satisfies (3.15), we find that $t^\pm(x^\pm)$

in Equation (3.15) should be given by,

$$t^\pm(x^\pm) = -\frac{N}{96(x^\pm)^2} . \quad (3.18)$$

Although $t^\pm(x^\pm)$ can be an arbitrary function of $x^\pm$ in general, we assume Equation (3.18) for a while. The contribution of the energy-momentum tensor of massless matter fields can effectively shift $t^\pm(x^\pm)$.

Equations (3.9), (3.15) and (3.17) can be easily solved. A static solution is given by,[★]

$$\begin{aligned} F_{+-} &= C J_0(p_0) \quad (x^- < 0) \\ &= C I_0(q_0) \quad (x^- > 0) , \\ e^{-2\phi} + \frac{N}{48}\phi &= \frac{M}{\lambda} - \lambda^2 x^+ x^- - \frac{N}{96} \ln(-\lambda^2 x^+ x^-) + \frac{1}{N} \left(\frac{2C}{g_A^2}\right)^2 L_0(x^+, x^-) \end{aligned} \quad (3.19)$$

Here $L_0(x^+, x^-)$ is defined by

$$\begin{aligned} L_0(x^+, x^-) &= -p_0^2 (J_0(p_0)^2 + 2J_1(p_0)^2 - J_0(p_0)J_2(p_0)) , \quad (x^- < 0) \\ &= q_0^2 (I_0(q_0)^2 - 2I_1(q_0)^2 + I_0(q_0)I_2(q_0)) , \quad (x^- > 0) \end{aligned} \quad (3.20)$$

and p_0 and q_0 are defined by,

$$p_0 \equiv \sqrt{-\frac{g_A^2 N x^- x^+}{2}}, \quad q_0 \equiv \sqrt{\frac{g_A^2 N x^- x^+}{2}} . \quad (3.21)$$

The solution (3.19) does not depend on “time” t when we change the coordinate by,

$$x^+ = e^{g(t)} f(r) , \quad x^- = e^{-g(t)} f(r) . \quad (3.22)$$

Here g and f are arbitrary functions.

★ There is an ambiguity for the definition of $\ln(-\lambda^2 x^+ x^-)$ when $\lambda^2 x^+ x^- > 0$.

We can also obtain the solutions describing the matter shock wave collapsing into the dilaton vacuum. When we consider the shock wave whose trajectory is $x^+ = x_0^+$, F_{+-} is given by Equation (3.11) and $e^{-2\phi} + \frac{N}{48}\phi$ is given by

$$e^{-2\phi} + \frac{N}{48}\phi = \frac{M}{\lambda} - \lambda^2 x^+ (x^- + \frac{M}{\lambda^3 x_0^+}) - \frac{N}{96} \ln(-\lambda^2 x^+ x^-) + \frac{1}{N} (\frac{2C}{g_A^2})^2 L(x^+, x^-) . \quad (3.23)$$

Here $L(x^+, x^-)$ is defined by

$$\begin{aligned} L(x^+, x^-) &= -p^2 (J_0(p)^2 + 2J_1(p)^2 - J_0(p)J_2(p)) , \quad (x^- < \tilde{x}_0) \\ &= q^2 (I_0(q)^2 - 2I_1(q)^2 + I_0(q)I_2(q)) . \quad (x^- > \tilde{x}_0) \end{aligned} \quad (3.24)$$

4. The Fate of Singularity and Apparent Horizon

In this section, we discuss the behavior of curvature singularity and apparent horizon by using the exact solution (3.23). In this section we assume $\tilde{\lambda}^2 \equiv \lambda^2 - \frac{2C^2}{g_A^2}$ is positive. The main results are following:

1. When $\lambda^3 x_0^+ \tilde{x}_0 > -M$, the essential behavior is similar to that of Callan, Giddings, Harvey and Strominger's dilatonic black hole theory modified in the paper by Russo, Susskind and Thorlacius.¹³ The singularity formed by the collapsing matter emerged naked.
2. When $\lambda^3 x_0^+ \tilde{x}_0 < -M$, the singularity and the apparent horizon disappear in a finite proper time. Furthermore, there appears a discontinuity in the metric on the trajectory of the collapsing matter ($x = x_0^+$, $x^- > -\frac{1}{\lambda^2 x_0^+} \frac{N}{96}$).

The curvature $R = 8e^{-2\rho} \partial_+ \partial_- \rho$ become singular when $e^{-2\phi} = \frac{N}{96}$ since

$$R \propto (e^{-2\phi} - \frac{N}{96})^{-1} . \quad (4.1)$$

By using Equation (3.23), we find the line of the singularity is given by,

$$\lambda^2 x^+ x^- + \frac{M}{\lambda x_0^+} (x^+ - x_0^+) + \frac{N}{96} \ln(-\lambda^2 x^+ x^-) + \frac{N}{96} (1 - \ln \frac{N}{96}) = \frac{1}{N} (\frac{2C}{g_A^2})^2 L(x^+, x^-) . \quad (4.2)$$

A point on the trajectory $(x^+, x^-) = (x_0^+, -\frac{1}{\lambda^2 x_0^+} \frac{N}{96})$ satisfies Equation (4.2).

We now discuss about the solution of the Equation (4.2) when x^+ is large enough. It is convenient to divide the both side of Equation (4.2) by x^+ and define functions $\mathcal{R}(x^-)$ and $\mathcal{L}(x^-)$ by

$$\begin{aligned}\mathcal{R}(x^-) &= \frac{1}{x^+} \left\{ \lambda^2 x^+ x^- + \frac{M}{\lambda x_0^+} (x^+ - x_0^+) + \frac{N}{96} \ln(-\lambda^2 x^+ x^-) + \frac{N}{96} (1 - \ln \frac{N}{96}) \right\}, \\ \mathcal{L}(x^-) &= \frac{1}{N x^+} \left(\frac{2C}{g_A^2} \right)^2 L(x^+, x^-).\end{aligned}\tag{4.3}$$

Then the singularity line is given by $\mathcal{R}(x^-) = \mathcal{L}(x^-)$. When $x^+ \rightarrow +\infty$, the curve $\mathcal{C}^{\mathcal{R}} = \{(x^-, y) | y = \mathcal{R}(x^-)\}$ approaches to the following curve

$$\begin{aligned}\mathcal{C}^{\mathcal{R}} \implies \mathcal{C}_{\infty}^{\mathcal{R}} &= \{(x^-, y) | y = \lambda^2(x^- + \frac{M}{\lambda^3 x_0^+}), \quad x^- < 0\} \\ &\oplus \{(x^-, y) | y < \frac{M}{\lambda x_0^+}, \quad x^- = 0\}\end{aligned}\tag{4.4}$$

On the other hand, the curve $\mathcal{C}^{\mathcal{L}} = \{(x^-, y) | y = \mathcal{L}(x^-)\}$ approaches to the following curve

$$\mathcal{C}^{\mathcal{L}} \implies \mathcal{C}_{\infty}^{\mathcal{L}} = \{(x^-, y) | y = 0, \quad x^- < \tilde{x}_0\} \oplus \{(x^-, y) | y \geq 0, \quad x^- = 0\}\tag{4.5}$$

When $\lambda^3 x_0^+ \tilde{x}_0 > -M$, there appear two solutions satisfying the equation $\mathcal{R}(x^-) = \mathcal{L}(x^-)$ in the limit $x^+ \rightarrow \infty$,

$$\begin{aligned}x^- &= -\frac{M}{\lambda^3 x_0^+}, \\ x^- &= 0 \quad (\tilde{x}_0 > 0) \quad \text{or} \quad x^- = \tilde{x}_0 \quad (\tilde{x}_0 < 0).\end{aligned}\tag{4.6}$$

Note that there appears two lines of singularities.

When $\lambda^3 x_0^+ \tilde{x}_0 < -M$, there is not any solution satisfying the equation $\mathcal{R}(x^-) = \mathcal{L}(x^-)$ when $x^+ = \infty$. This tells that the curvature singularities disappear in a finite proper time. The disappearance occurs in a weak coupling region

$e^{-2\phi} < \frac{N}{96}$ and the semi-classical analysis given here is reliable. The collapsing matter ‘pair creates’ two curvature singularities and these singularities ‘pair annihilate’ in a finite proper time when $\lambda^3 x_0^+ \tilde{x}_0 < -M$, *i.e.*, the line of the singularity makes a loop in x^+x^- plane. Outside of the loop, we find $e^{-2\phi} < \frac{N}{96}$ in a region $x^+ > x_0^+$. On the other side, in a region $x^+ < x_0^+$ where the solution describes the dilaton vacuum (2.10), $e^{-2\phi} > \frac{N}{96}$ just below the trajectory of collapsing matter $x^+ = x_0^+ - 0$ when $x^- > -\frac{1}{\lambda^2 x_0^+} \frac{N}{96}$. Therefore there appears a discontinuity on the trajectory $x^+ = x_0^+$, $x^- > -\frac{1}{\lambda^2 x_0^+} \frac{N}{96}$ although $e^{-2\phi} + \frac{N}{48}\phi$ is continuous. The discontinuity would be a signal of topology change.

Although the curvature singularity appears at $(x^+, x^-) = (x_0^+, -\frac{1}{\lambda^2 x_0^+} \frac{N}{96})$, the singularity disappears on the line $x^+ = x_0^+ + \epsilon$ (ϵ is an infinitesimal constant) if

$$\partial_+ \mathcal{R}(x^-)|_{x^+=x_0^+} < \partial_+ \mathcal{L}(x^-)|_{x^+=x_0^+} , \quad (4.7)$$

that is,

$$\tilde{x}_0 < -\frac{Mg_A^2}{2C^2\lambda x_0^+} - \frac{N}{96\lambda^2 x_0^+} . \quad (4.8)$$

In this case, the curvature singularities appear after a finite proper time or any singularity does not appear. In any case, the discontinuity of the metric appears on the trajectory of the collapsing matter.

Since $e^{-2\phi} < \frac{N}{96}$ outside the singularity loop in a region $x^+ > x_0^+$, $e^{-2\phi} \sim -\frac{N}{96} \ln(-\lambda^2 x^+ x^-)$ when $-\lambda^2 x^+ x^- \sim 0$. Then the curvature $R = 8e^{-2\rho} \partial_+ \partial_- \rho$ diverges as $R \sim \{-\lambda^2 x^+ x^- \ln(-\lambda^2 x^+ x^-)\}^{-1}$. Therefore there appears a new line of the singularity when $x^- = 0$. The singularity occurs since we have chosen $t^\pm(x^\pm)$ by Equation (3.18). There is an ambiguity to how to choose $t^\pm(x^\pm)$ since $t^\pm(x^\pm)$ or $r^\pm(x^\pm)$ in Equation (3.16) can be regarded as a part of dynamical modes of the field Z in Equation (3.14). If we choose $t^-(x^-)$, for example, by $t^-(x^-) = -\frac{N}{96} \frac{1}{2} \partial_-^2 \ln\{(x^-)^2 + \alpha^2\}$ (α is a constant), there is no singularity except the singularities coming from $e^{-2\phi} = \frac{N}{96}$. Therefore the singularity which appeared when $-\lambda^2 x^+ x^- \sim 0$ would not have a physical meanings. If we do not change the

asymptotic behavior of $t^\pm(x^\pm)$, the qualitative behavior of the $e^{-2\phi} = \frac{N}{96}$ singularity line do not depend on the choice of $t^\pm(x^\pm)$.

We now discuss about the apparent horizon which is defined by, $\partial_+ \phi = 0$.⁷ By using Equation (3.23), we find that the apparent horizon is given by,

$$\lambda^2(x^- + \frac{M}{\lambda^3 x_0^+}) + \frac{N}{96x^+} = \frac{1}{N}(\frac{2C}{g_A^2})^2 \partial_+ L(x^+, x^-) . \quad (4.9)$$

When $x^+ = x_0^+ + 0$, the solutions which satisfies Equation (4.9) is given by, $x^- = -\frac{1}{\lambda^2}(\frac{M}{\lambda x_0^+} + \frac{N}{96x_0^+} + \frac{2C^2 \tilde{x}_0}{g_A^2})$ and $x^- = -\infty$. Note that there appears two lines of apparent horizons in general. We define $\mathcal{R}^A(x^-)$ and $\mathcal{L}^A(x^-)$ by

$$\begin{aligned} \mathcal{R}^A(x^-) &= \lambda^2(x^- + \frac{M}{\lambda^3 x_0^+}) + \frac{N}{96x^+} , \\ \mathcal{L}^A(x^-) &= \frac{1}{N}(\frac{2C}{g_A^2})^2 \partial_+ L(x^+, x^-) . \end{aligned} \quad (4.10)$$

The apparent horizon is given by $\mathcal{R}^A(x^-) = \mathcal{L}^A(x^-)$. When $x^+ \rightarrow +\infty$, the curves $\mathcal{C}^{AR} = \{(x^-, y) \mid y = \mathcal{R}^A(x^-)\}$ and $\mathcal{C}^{AL} = \{(x^-, y) \mid y = \mathcal{L}^A(x^-)\}$ approach to the following curves

$$\begin{aligned} \mathcal{C}^{AR} &\implies \mathcal{C}_\infty^{AR} = \{(x^-, y) \mid y = \lambda^2(x^- + \frac{M}{\lambda^3 x_0^+})\} \\ \mathcal{C}^{AL} &\implies \mathcal{C}_\infty^{AL} = \{(x^-, y) \mid y = 0, \quad x^- < \tilde{x}_0\} \oplus \{(x^-, y) \mid y \geq 0, \quad x^- = 0\} \end{aligned} \quad (4.11)$$

When $\lambda^3 x_0^+ \tilde{x}_0 > -M$, there appear two solutions satisfying the equation $\mathcal{R}^A(x^-) = \mathcal{L}^A(x^-)$ in the limit of $x^+ = \infty$,

$$x^- = -M\lambda^3 x_0^+ , \quad x^- = \tilde{x}_0 . \quad (4.12)$$

In this case, the behaviors of the lines of singularities and apparent horizons are qualitatively similar to that of Callan, Giddings, Harvey and Strominger's dilatonic black hole theory modified in the paper by Russo, Susskind and Thorlacius¹³ and the singularity emerges naked. When $\lambda^3 x_0^+ \tilde{x}_0 < -M$, there is not any solution satisfying the equation $\mathcal{R}^A(x^-) = \mathcal{L}^A(x^-)$ in the limit of $x^+ = \infty$ and the apparent horizons disappear in a finite proper time.

5. Summary and Discussion

We have analyzed exactly solvable semi-classical theory of two dimensional dilatonic gravity with electromagnetic interactions. We added the term which changes the kinetic term and keeps the current $J^\mu = \partial^\mu(\phi - \rho)$ to conserve. The theory contains massless fermions as matter fields and there appear the quantum corrections including chiral anomaly. The screening effect due to the chiral anomaly has a tendency to cloak the singularity. When $\lambda^3 x_0^+ \tilde{x}_0 < -M$, the singularity coming from $e^{-2\phi} = \frac{N}{96}$ and the apparent horizon disappear in a finite proper time. Furthermore, there appears a discontinuity in the metric on the trajectory of the collapsing matter, which would be a signal of topology change.

One of the problems is whether the disappearance of the singularities observed here is independent of the details of the models since many semi-classical models have been proposed.^{10–13} We have also analyzed the model without ϕR term in (1.1), which corresponding to the original model by CGHS. By assuming that $\tilde{\lambda}^2 \equiv \lambda^2 - \frac{2C^2}{g_A^2}$ is small compared to λ^2 , we have investigated the line $x^- = f(x^+)$ where ϕ is a constant (on the singularity $\phi = -\frac{1}{2}\ln \frac{N}{96}$) and we find

$$f(x^+) = f_0 - f_1 \frac{\tilde{\lambda}^2}{\lambda^2} (x^+ - x_0^+) + f_2 (x^+ - x_0^+)^2 + \mathcal{O}((x^+ - x_0^+)^3). \quad (5.1)$$

Here f_i 's are order one quantities with respect to $\frac{\tilde{\lambda}^2}{\lambda^2}$ and positive if $\tilde{x}_0 \ll -\frac{N}{24\lambda^2 x_0^+}$. Therefore there exists a point $x^+ = \tilde{x}_0^+$ which satisfies $\tilde{x}_0^+ - x_0^+ = \mathcal{O}(\frac{\tilde{\lambda}^2}{\lambda^2})$, $\tilde{x}_0^+ - x_0^+ > 0$ and $f'(x^+) = 0$. This tells that there is a turning point and we have found that the screening effect due to the chiral anomaly has a tendency to cloak the singularity.

Acknowledgements: We acknowledge K. Odaka and A. Sugamoto for discussions. We are also indebted to S. Odake and T. Tada for useful informations. The research of I.O. is supported by the Japan Society for the Promotion of Science.

REFERENCES

1. S.W. Hawking, *Comm.Math. Phys.* **43** (1975), 199
2. G. 't Hooft, *Nucl.Phys.* **B335** (1990), 138
3. Y. Aharonov, A. Casher and S. Nissimov, *Phys.Lett.* **B191** (1987), 51
4. C.F.E. Holzhey and F. Wilczek, preprint IASSNS-HEP-91/71
5. C.G. Callan, S.B. Giddings, J.A. Harvey and A. Strominger, *Phys.Rev.* **D45** (1992), R1005
6. J.G. Russo, L. Susskind and L. Thorlacius, preprint SU-ITP-92-4 (1992)
7. L. Susskind and L. Thorlacius, preprint SU-ITP-92-12 (1992)
8. S.W. Hawking, preprint CALT-68-1774
9. T. Banks, A. Dabholkar, M.R. Douglas and M. O'Loughlin, *Phys.Rev.* **D45** (1992), 3607
10. A. Bilal and C.G. Callan, preprint PUPT-1320 (1992)
11. S.P. de Alwis, preprints, COLO-HEP-280, 284 (1992)
12. K. Hamada, preprint UT-Komaba 92-7
13. J.G. Russo, L. Susskind and L. Thorlacius, preprint SU-ITP-92-17 (1992)
14. S. Nojiri and I. Oda, preprint NDA-FP-5/92, OCHA-PP-26 (1992)
15. N. Ishibashi, M. Li and R. Steif, *Phys.Rev.Lett.* **67** (1991), 3336
16. M.D. McGuigan, C.R. Nappi and S.A. Yost, *Nucl.Phys.* **B375** (1992), 421